

09/949784

(10)



Europäisches Patentamt

European Patent Office

Office européen des brevets

(11) Publication number:

0 109 561

A1

Party
#9

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 83110477.3

(51) Int. Cl.³: A 61 K 31/17

(22) Date of filing: 20.10.83

(30) Priority: 27.10.82 US 437072

(63) Date of publication of application:
30.05.84 Bulletin 84/22

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: USV PHARMACEUTICAL CORPORATION
1 Scarsdale Road
Tuckahoe New York(US)

(72) Inventor: Wolf, Peter S.
Richard Somers Road
Granite Springs New York(US)

(72) Inventor: Kesselman, Morris A.
21 Autumn Circle
Yonkers New York(US)

(74) Representative: Patentanwälte Grünecker, Dr.
Kinkeldey, Dr. Steckmair, Dr. Schumann, Jakob, Dr.
Bezold, Meister, Hilgers, Dr. Meyer-Pfath
Maximilianstrasse 58
D-8000 München 22(DE)

(54) Celiprolol for the treatment of glaucoma.

(57) An ophthalmic preparation for the treatment of glaucoma by the application to the glaucomatous eye of a celiprolol salt in a pharmaceutically acceptable ophthalmic carrier.

EP 0 109 561 A1

1 CELIPROLOL FOR THE TREATMENT OF GLAUCOMA

 This invention relates to compositions and methods
for treating intraocular pressure associated with glaucoma.
More particularly it relates to the use of celiprolol
5 hydrochloride, as well as selected pharmaceutically
acceptable salts thereof, that have been found useful in
lowering intraocular pressure.

 Elevated intraocular pressure is a major risk
factor in the onset and development of glaucomatous visual
10 field loss. The higher the level of intraocular pressure,
the greater the likelihood of glaucomatous visual field loss
and optic nerve damage.

 Attempts have been made to lower intraocular
pressure in glaucoma by administering the patients certain
15 beta adrenergic blocking agents, commonly known as beta
blockers.

 In the human body, the beta blockers have several
effects. For example, they can reduce the heart rate of an
angina patient, which in turn reduces the workload of the
20 heart and thus its need for blood and oxygen. They also tend
to decrease the heart's force of contraction, which likewise
diminishes the heart's workload. In addition, these drugs
reduce the systolic blood pressure which is beneficial to
patients with hypertension.

25 In addition to treating heart ailments, some beta
blockers were experimented with for treating other ailments,
such as migraine, alcohol and drug withdrawal problems, and
glaucoma.

30 While the use of beta blockers provides valuable
benefits to humankind, it also has some undesirable side
effects and reactions. Some beta blockers tend to build up
in the central nervous system causing fatigue, lethargy, and

1 confusion. Others may cause bronchial spasm and cannot be
used in people with bronchial asthma. Still others, while
having minimal side effects, fail to produce acceptable
results because of their limited potency.

5 Timolol, a non-selective beta-adrenoceptor
antagonist, has been shown to lower intraocular pressure in
both patients with normal intraocular pressure and in
patients with open angle glaucoma. Sold as TIMOPTIC[®], it
is the only beta block sold in the U.S. for this purpose.
10 Among the adverse reactions reported on the use of timolol is
the aggravation or precipitation of certain cardiovascular
and pulmonary disorders including bronchospastic disease,
sinus bradycardia, cardiogenic shock and cardiac failure.
Similarly, atenolol, sotalol, pindolol, oxprenolol,
15 practolol, propranolol, butidrine and metoprolol were
reported to have activity in the treatment of glaucoma.
However, some of these compounds have been reported to cause
pronounced side-effects, for example, metoprolol provokes
allergic reactions, oxprenolol may cause corneal
20 epitheliopathy and practolol induces oculomucoculaneous
syndrome by immunopathological reactions.


Celiprolol hydrochloride has been shown a selective
beta-1-adrenoceptor antagonist having intrinsic
sympathomimetic, but no local anesthetic activity.

25 It has now been discovered that celiprolol
hydrochloride, as well as selected pharmaceutically
acceptable salts thereof such as maleate, succinate and the
like, when topically applied to the eyes in a
pharmaceutically suitable vehicle, such as an ophthalmic
30 solution, is effective in lowering intraocular pressure.
Such ophthalmic preparation contain from about 0.01% w/v to
about 5.0% w/v, preferably from about 0.03 w/v to about 2.0%

w/v of the active ingredient along with inactive ingredients used in the art, such as sodium borate-boric acid, sodium hydroxide to adjust pH, benzalkonium chloride as preservative and water as the vehicle of preference.

It has also been surprisingly discovered that celiprolol hydrochloride, when used according to the present invention, produces bronchodilation following each application and as such possesses significant therapeutic advantage to patients suffering from both glaucoma and respiratory disease.

Celiprolol hydrochloride (3-[3-acetyl-4-[3(tert-butylamino)-2-hydroxy-propoxy]-phenyl]-1, 1-diethylurea hydrochloride) as the free base has the following structure:



It has a molecular weight of 415.95 having C = 57.75%, H = 8.24%, N = 10.10%, O = 15.39% and Cl = 8.52%.

Celiprolol can be prepared according to several pathways described in Austrian Patents: 334,385 issued to Zoelss, G. et al.; 335,465 to Zoelss, G.; 335,464 to Zoelss, G.; and 335,467 to Zoelss, G. The pathways for preparation of celiprolol according to these patents are as illustrated:

1 The inactive carriers for the active compounds used
in the formulations of the present invention include water
and ointment bases, such as mineral oil in the range of about
2 to about 10% w/v and white petrolatum in the range of about
5 90 to 98% w/v. In preparing the formulations of the present
invention, the active compound is solubilized in the carrier.
For solubilizing the active compound a co-solvent, in
addition to the carrier, may be used. Such co-solvents
include glycerin polyethylene glycol fatty acid esters in the
10 range of 1 to 10% w/v, propylene glycol in the range of 1 to
10% w/v, polyethylene glycol in the range of 1 to 15% w/v,
polysorbate 20, 60 and 80 in the range of 0.01 to 0.2% w/v
and Pluronic F-68 in the range of 0.01 to 2% w/v and mixtures
thereof.

15 To prevent irritation to the eye the isotonicity of
the preparation should be in the range of 270 to 330
milliosmoles. Sodium chloride in the range of $0.9 \pm 0.1\%$ w/v
may be used, if necessary, to adjust the isotonicity.

20 The ophthalmic preparations of the present
invention will have a pH of about 6 to 9 preferably in the
range of 7-8. Buffers that may be used to obtain said pH
range include alkali metal or alkaline metal earth
carbonates, bicarbonates, borates, citrates and tris buffers.
More specifically such buffers include 0.01 to 0.2 molar
25 concentrations of boric acid-sodium borate, phosphate
buffers, boric acid-sodium bicarbonate, boric acid-sodium
citrate, citric acid-sodium phosphate, tris(hydroxymethyl)
amino methane-maleic acid and tris(hydroxymethyl) amino
methane-HCl.

30 Other ingredients which may be desirable to use in
the ophthalmic preparations of the present invention include
viscosity builder agents, preservatives and stabilizers.

1 Examples of these, which may be incorporated into the preparations during the process or after the active compound is solubilized, include the following:

- 5 a. preservatives in the range of about 0.001 to about 1.0% w/v;
- b. stabilizers in the range of about 0.01 to about 5.0% w/v; and viscosity builders or viscosity agents in the range of about 0.01 to about 2.0% w/v.

More specific examples include:

10	<u>Preservatives</u>	<u>Range % w/v</u>
	Benzalkonium Chloride	0.004 - 0.02
	Disodium Ethylenediamine Tetraacetate	0.01 - 0.20
	Thimerosal	0.001 - 0.01
	Chlorobutanol	0.5 - 1.0
15	Phenylmercuric nitrate	0.002 - 0.02
	Phenylmercuric acetate	0.002 - 0.02
	Methyl Paraben	0.03 - 0.20
	Propyl Paraben	0.01 - 0.05
	Phenylethyl alcohol	0.25 - 0.75
	Phenyl mercuric borate	0.002 - 0.02
20	<u>Stabilizers</u>	<u>Range % w/v</u>
	Sodium Bisulfite	up to 0.5%
	Sodium Thiosulfite	up to 0.5%
	Cysteine	up to 3%
	Acetyl cysteine	up to 3%
	B-cyclodextrin	up to 3%
25	Dextran	up to 5%
	Thiourea	up to 3%
	Thiosorbitol	up to 3%
	Monothioglycerol disodium EDTA	up to 3%
	Dioctyl Sodium Sulfosuccinate	up to 3%
30	<u>Viscosity Agents</u>	<u>Range % w/v</u>
	Polyvinylpyrrolidone	0.5 - 2.0%
	Polyvinyl alcohol	0.5 - 2.0%
	Methyl cellulose	0.1 - 1.0%
	Hydroxypropyl Methylcellulose	0.10 - 1.0%
	Hydroxyethyl cellulose	0.10 - 1.0%
35	Carboxymethyl cellulose	0.10 - 1.0%
	Sodium Carboxymethyl cellulose	0.01 - 1.0%
	Hydroxypropylcellulose	0.01 - 1.0%

In general, the preparations of the present invention may be made and manufactured as illustrated herein;

a., The active drug is dissolved in the aqueous vehicle or thoroughly dispersed in the ointment vehicle by adequate stirring;

b., After dissolution or dispersion of the active drug additional ingredients, such as preservatives, buffer salts, stabilizers and viscosity agents are added and dissolved by further stirring. Sodium chloride is then added, if required, to adjust isotonicity, and the solution is brought to final volume;

c., The product is then sterilized by filtration through a 0.22 micron membrane or alternatively autoclaved at 121-123°C, or by a combination of both methods;

d., The sterile solution is filled into sterile containers and sealed.

The following examples illustrate the present invention without, however, limiting the same thereto.

EXAMPLE A

	<u>% w/v</u>
Celiprolol	0.50
Benzalkonium Chloride	0.01
DiSodium EDTA	0.10
Purified Water, Q.S.	100.00

EXAMPLE B

	<u>% w/v</u>
Celiprolol	0.50
Benzalkonium Chloride	0.01
DiSodium EDTA	0.10
Boric Acid	0.215
Sodium Borate, Q.S. to pH 7.4	
Purified Water, Q.S.	100.00

1

EXAMPLE C% w/v

5

Celiprolol	0.125
Thimerosal	0.002
Tris (Trihydroxyl methyl amino methane	0.12
Maleic Acid, Q.S. to pH 8.3	
Purified Water, Q.S.	100.00

10

EXAMPLE D% w/v

15

Celiprolol	1.00
Benzalkonium Chloride	0.01
Boric Acid	1.115
Sodium Borate, Q.S. to pH 7.4	
Polyvinyl Alcohol	1.4
Purified Water, Q.S.	100.00

EXAMPLE E% w/v

20

Celiprolol	3.00
Chlorobutanol	0.50
Boric Acid	0.05
Sodium Thiosulfate	0.30
Sodium Bicarbonate, Q.S. to pH 7.0	
Purified Water, Q.S.	100.00

EXAMPLE F% w/v

25

30

Celiprolol	5.00
Phenyl Mercuric Acetate	0.02
Methyl Paraben	0.05
Propyl Paraben	0.01
Polyvinyl Pyrrolidone	2.00
Polysorbate 80	0.05
Purified Water, Q.S.	100.00

35

1

EXAMPLE G

5

	<u>% w/v</u>
Celiprolol	0.50
Benzalkonium Chloride	0.02
DiSodium EDTA	0.10
Boric Acid	0.05
Sodium Borate, Q.S. to pH 6.0	
Polyvinyl Pyrrolidone	1.50
Hydroxyethyl Cellulose	0.20
Purified Water, Q.S.	100.00

10

EXAMPLE H

15

	<u>% w/v</u>
Celiprolol	0.50
Benzalkonium Chloride	0.005
DiSodium EDTA	0.05
Boric Acid	0.215
Sodium Borate to adjust pH to 7.4	
Polyethylene Glycol 300	5.0
Glycerin	1.0
Sodium Thiosulfate	0.5
Purified Water, Q.S.	100.00

20

EXAMPLE I

25

	<u>% w/v</u>
Celiprolol	1.0
Methyl Paraben	0.05
Propyl Paraben	0.01
Mineral Oil	5.0
White Petrolatum, Q.S.	100.00

EXAMPLE J

30

	<u>% w/v</u>
Celiprolol	0.5
Chlorobutanol	0.5
Polyethylene Glycol 400	4.0
White Petrolatum, Q.S.	100.00

35

1 Illustrative of the benefits obtained in accordance
with the present invention the studies described in the
Examples following, were conducted.

EXAMPLE 1

5 This example shows that celiprolol decreases
intraocular pressure in dogs and that the decrease is dose-
dependent.

Animals, Testing Procedure and Apparatus

10 Mongrel dogs of either sex weighing between 9 and
16 kg were anesthetized with sodium pentobarbital, 35 mg/kg,
i.v. (Ganes Chemical, Pennsville, N.J.). The animals then
were intubated with an endotracheal tube (Rusch, size 8-9F,
Artistic Surgical, New York, N.Y.) and allowed to breathe
spontaneously.

15 Pulsatile arterial pressure was monitored using a
polyethylene catheter (PE 240, Clay-Adams, Parsippany, N.J.)
inserted into the right femoral artery and its tip advanced
until a distinct dicrotic notch was observed on the arterial
pressure tracing. The catheter was connected to a pressure
20 transducer (P23 ID, Statham, Oxnard, CA) and a D.C. driver
amplifier (Model 7D; Grass Instruments, Quincy, MA) via a low
leve D.C. preamplifier (Model 7Pl, Grass Instruments). Mean
arterial pressure was determined electronically by damping
the pulsatile arterial pressure signal. Heart rate was
25 recorded from the output of a tachometer (Model 7PA, Grass
Instruments) triggered by the R wave of a Lead II
electrocardiogram (EKG -Tachograph Preamplifier, Model 7P4,
Grass Instruments). The outputs of arterial pressure and EKG
-Tachometer Preamplifier were recorded continuously on an
30 oscillograph (Model 7D, Grass Instruments). Heart rate was
measured manually from the tachometer output or calculated
from the EKG preamplifier output. Mean arterial pressure was
measured manually from the oscillograph tracings.

1 Intraocular pressure was measured using a pneumatonometer
(Model 30R, Digilab, Cambridge, MA).

5 Upon completion of all surgical procedure the
animals were allowed to stabilize for 15-30 minutes before
pretreatment baseline measurements were taken.

PROTOCOL

10 Eight mongrel dogs were divided into two groups of
four dogs each: the control group and the test group. Two
pretreatment baseline measurements of intraocular pressure,
mean arterial pressure and heart rate were recorded at -15
and 0 minutes. The means values of these two readings were
used as the pretreatment (0 time) values. The control group
was then administered saline topically in a volume of 5 μ l
instilled into the left eye of each dog at hourly intervals.
15 The test group was administered celiprolol topically in a
volume to 50 μ l instilled into the left eye of each dog, in
ascending concentrations (0.03%, 0.06%, 0.125%, 0.5%), at
hourly intervals. Intraocular pressure, mean arterial
pressure and heart rate were recorded at 15 minute intervals
20 for 60 minutes post drug or vehicle (saline) administration
for each concentration of celiprolol administered and each
administration of saline.

Drug Preparation

25 Celiprolol (RHC 5320-A Lot 2) was dissolved and
diluted in normal saline (0.9%, Abbott Laboratories, North
Chicago, IL).

The control group received normal saline.

Data Analysis

30 The absolute change and the percent change from the
pretreatment (0 time) values for intraocular pressure, mean
arterial pressure and heart rate were expressed as the mean \pm
S.D. The significance of the difference between the vehicle

1 control group and the celiprolol test group were evaluated
using a "t" test for grouped data (as described in SAS T TEST
Release 79.5, SAS Institute, Gary, NC 1982). Differences
were considered significant if P 0.05. In addition, the
5 maximum percent intraocular pressure change for each
concentration of celiprolol administered was expressed as the
mean \pm 1 S.D.

Result

Test results were shown in Tables I, II & III.

10

15

20

25

30

35

TABLE I
THE EFFECT OF ASCENDING CONCENTRATIONS OF CELIPROLOL ON INTRAOCULAR PRESSURE (IOP) IN ANESTHETIZED DOGS

Treatment	N	Baseline ^a Pre-treatment (mm Hg)	0.03%					0.06%					0.125%					0.25%					0.5%				
			15	30	45	60	75	90	105	120	135	150	165	180	195	210	225	240	255	270	285	300					
Saline (50 µl of 0.9% NaCl Solution)	4	21 +2	0 +2	-1 +3	-2 +2	-3 +2	-2 +3	-2 +3	-4 +3	-4 +2	-4 +2	-4 +2	-5 +1	-5 +3	-5 +2	-6 +2	-5 +1	-5 +1	-5 +4	-4 +3	-5 +1	-4 +1	Z Change	-4 +5	-5 +4	-20 +13	-25 +6
			+10	+13	+10	+8	+11	+8	+13	+12	+11	+9	+7	+16	+8	+8	+5	+3	+16	+13	+6	+4					
Celiprolol HCl (RUC 5320-A-2 50 µl)	4	30 +8	-4 +5	-5 +4	-5 +3	-9 +7	-10 ^b +5	-10 ^b +5	-11 ^b +5	-10 ^b +2	-11 ^b +4	-13 ^b +6	-11 +6	-11 +6	-13 ^b +6	-12 +7	-14 ^b +6	-8 +4	-12 +8	-20 ^b +7	-14 +7	-15 ^b +6	Z Change	-11 +15	-28 +19	-41 +20	-46 ^b +12
			+10	+13	+10	+8	+11	+8	+13	+12	+11	+9	+7	+16	+8	+8	+5	+3	+16	+13	+6	+4					

^aAll values are the mean \pm 1 S.D.

^bp < 0.05

TABLE II
THE EFFECT OF ASCENDING CONCENTRATIONS OF CELIPROLOL ON MEAN ARTERIAL PRESSURE (MAP) IN ANESTHETIZED DOGS

Treatment	N	Baseline ^a Pre-treatment (mm Hg)	Dose Time Minutes					0.06%	0.1%					0.3%					0.5%											
			15	30	45	60	75		90	105	120	135	150	165	180	195	210	225	240	255	270	285	300							
Saline (50 µl of 0.9% NaCl Solution)	4	113 ±10	8 +5	6 ±15	4 ±11	8 ±10	9 ±18	9 ±8	9 ±7	10 ±12	6 ±15	11 ±13	8 ±16	7 ±17	12 ±14	6 ±10	6 ±9	7 ±8	7 ±6	1 ±13	3 ±11	7 ±11								
	2	Change	7 +4	5 ±13	3 ±10	7 ±9	8 ±16	8 ±7	8 ±6	8 ±10	5 ±13	10 ±12	7 ±14	6 ±15	11 ±13	5 ±9	6 ±8	7 ±7	6 ±5	0 ±11	3 ±10	6 ±9								
Celiprolol HCl (RUC 5320-A-2 50 µl)	4	126 ±15	-1 +10	2 ±15	-3 ±9	-3 ±8	-7 ±7	-7 ±10	-8 ^b ±8	-7 ±9	-7 ±10	-3 ±9	-2 ±5	11 ±22	-4 ±10	-3 ±2	-4 ±7	-5 ±16	-5 ±16	-4 ±11	-6 ±14	-9 ±20								
	2	Change	0 +8	2 ±11	-2 ±7	-2 ±6	-5 ±5	-6 ^b ±8	-5 ^b ±6	-6 ±7	-6 ±8	-2 ^b ±7	-1 ±4	10 ±19	-4 ±9	-2 ±2	-3 ±6	-5 ±14	-4 ±14	-3 ±9	-5 ±12	-7 ±17								

^aAll values are the mean ± 1 S.D.
^bp < 0.05

TABLE III
THE EFFECT OF ASCENDING CONCENTRATIONS OF CELIPROLOL ON HEART RATE (BPM) IN ANESTHETIZED DOGS

Treatment	N	Baseline ^a Pre-treatment (mm Hg)	Dose time Minutes Post-treatment	0.03%					0.06%					0.1%					0.3%					0.5%				
				15	30	45	60	75	90	105	120	135	150	165	180	195	210	225	240	255	270	285	300					
Saline (50 µl of 0.9% NaCl Solution)	4	153 ±15	Change (bpm)	-9 +9	-12 +11	-17 +14	-17 +18	-21 +17	-18 +21	-14 +19	-9 +26	-24 +35	-21 +31	-17 +28	-23 +35	-24 +35	-32 +22	-26 +22	-6 +20	-23 +14	-30 +29	-27 +14	-26 +33	x Change	-14 +22	-17 +25	-18 +29	-13 +22
				-6 +5	-8 +7	-11 +9	-11 +12	-15 +11	-12 +14	-9 +13	-6 +17	-16 +23	-14 +21	-11 +18	-15 +22	-16 +23	-21 +15	-17 +15	-4 +13	-15 +10	-20 +19	-18 +10	-14 +22					
Celliprolol HCl (RHC 5320-A-2) 50 µl)	4	135 ±14	Change (bpm)	0 +4	-6 +9	-5 +12	-5 +14	-8 +14	-8 +13	-9 +26	-10 +16	-15 +22	-5 +33	-9 +33	6 +26	-3 +29	-11 +17	-17 +13	-17 +18	-11 +25	-8 +18	-18 +29	-12 +35	x Change	-14 +22	-17 +25	-18 +29	-13 +22
				0 +3	-4 +7	-3 +9	-4 +11	-6 +11	-5 +11	-7 +20	-7 +12	-11 +16	-3 +25	-6 +24	5 +20	-1 +23	-7 +13	-12 +8	-12 +13	-7 +18	-5 +13	-13 +22	-14 +26					

^a All values are the mean ± 1 S.D.

^b P < 0.05

1 Table I shows that celiprolol caused an immediate
decrease in intraocular pressure which persisted for the
duration of the experiment. The magnitude of the decrease
became significantly different ($P < 0.05$) from that of the
5 control group at 75 minutes. The data also indicate that
celiprolol causes a dose-dependent decrease in intraocular
pressure which reaches a maximum at 0.25% concentration.

 Table II shows that the celiprolol group exhibited
a small decrease in mean arterial pressure which became
10 significantly different from that of the control at a
concentration of 0.06%. Since this decrease in arterial
pressure was not dose-dependent, it may be a chance occurrence
and not pharmacologically important.

 Table III shows that celiprolol had no effect on
15 heart rate at any of the concentrations tested.

20

25

30

35

EXAMPLE II

This example compares the activity of celiprolol and timolol in dogs on intraocular pressure, systemic arterial pressure and heart rate.

Animals, Testing Procedure and Apparatus

Same as described in Example I.

Drug Preparation

Celiprolol and saline were used as shown in Example I. Timolol (TIMOPTIC[®], 0.5%, Merck Sharp and Dohme, West Point, PA) was used.

PROTOCOL

Mongrel dogs were screened for intraocular pressure and only dogs that had intraocular pressures between 23 mmHg and 28 mmHg were chosen in this study.

Twelve mongrel dogs which met the above-described criterion were subdivided into three groups of four dogs each. The control group had a mean intraocular pressure of 25.3 ± 0.6 mmHg; the celiprolol, or test, group had a mean intraocular pressure of 25.6 ± 0.6 mmHg; and the timolol group, with which the celiprolol group was compared, had a mean intraocular pressure of 25.2 ± 1.6 mmHg.

Two pretreatment baseline measurements of intraocular pressure, mean arterial pressure and heart rate were recorded at -15 and 0 minutes for each groups. The mean values of these two readings were used as pretreatment (0 time) values. 0.5% celiprolol in a volume of 50 μ l was then administered into the left of each dog in the test or celiprolol group; 0.5% timolol in a volume of 50 μ l was administered into the left eye of each dog in the timolol group; and 50 μ l of saline was administered into the left eye of each dog in the control group. Intraocular pressure, mean arterial pressure and heart rate measurements were made at various intervals for five hours respectively for post drug or saline administration.

1 Date Analysis

Same as in Example 1.

Results

5 Test results are shown in Tables IV, V and VI Table
IV shows that at equal doses both celiprolol and timolol were
effective in decreasing intraocular pressure in anesthetized
dogs. Celiprolol, however, caused a significant change from
control of the one hour post drug administration which
persisted for the duration of the study, while timolol
10 attained a significant difference from the control only after
180 minutes post drug administration, which from thereon also
persisted for the duration of the experiment.

Table V shows that neither celiprolol nor timolol
had any significant effect on means arterial pressure.

15 Table VI shows that celiprolol seems to have no
effect on heart rate whereas timolol at least for the first
two hours has a tendency to decrease heart rate.

20

25

30

35

TABLE IV
THE EFFECT OF TOPICALLY ADMINISTERED CELIPROLOL AND TIMOLOL ON INTRAOCULAR PRESSURE (IOP) IN ANESTHETIZED DOGS

Treatment	N	Baseline ^a Pre-treatment (mm Hg)	Time (Minutes Posttreatment)											
			15	30	45	60	90	120	150	180	210	240	270	300
Saline 50 μ l (0.9% NaCl Solution)	4	25 +1	-1 +3	-2 +5	0 +4	0 +3	0 +2	-2 +3	-1 +5	0 +4	0 +4	1 +5	1 +2	-2 +2
		Change (mm Hg)												
Celliprolol HCl (RMC 5320-A) 50 μ l 0.5% Solution	4	25 +1	-5 +1	-6 +2	1 +1	-1 +1	-1 +9	-8 +10	-5 +19	-1 +16	-1 +15	-4 +19	3 +7	-6 +10
		Change (mm Hg)												
Timolol Maleate (Timoptic) 50 μ l of 0.5 Solution	4	25 +2	-15 +0	-11 +5	-16 +10	-23 +6	-21 +6	-25 +8	-33 +11	-31 +7	-37 +9	-36 +7	-38 +5	-41 +8
		Change (mm Hg)												
		Change (mm Hg)												
		Change (mm Hg)												

^a All values are expressed as the mean \pm 1 S.D.
bp < 0.05

TABLE V
THE EFFECT OF TOPICALLY ADMINISTERED CELIPROLOL AND TIMOLOL ON MEAN ARTERIAL PRESSURE (MAP) IN ANESTHETIZED DOGS

Treatment	N	Baseline ^a Pre-treatment (mm Hg)	Time (Minutes Posttreatment)											
			15	30	45	60	90	120	150	180	210	240	270	300
Saline 50 μ l (0.9% NaCl Solution)	4	120 \pm 13	4	3	3	2	2	-2	2	5	5	2	0	5
			\pm 3	\pm 8	\pm 4	\pm 6	\pm 5	\pm 11	\pm 10	\pm 5	\pm 13	\pm 11	\pm 8	\pm 9
			Change											
Celiprolol HCl (RHC 5320-A) 50 μ l 0.5% Solution	4	113 \pm 8	3	2	3	2	2	-2	2	4	4	2	0	4
			\pm 2	\pm 6	\pm 3	\pm 5	\pm 5	\pm 10	\pm 9	\pm 5	\pm 12	\pm 10	\pm 7	\pm 8
			Change											
Timolol Maleate (Timoptic) 59 μ l of 0.5% Solution	4	106 \pm 6	2	4	3	4	5	10	0	3	4	2	-2	-1
			\pm 2	\pm 3	\pm 6	\pm 7	\pm 8	\pm 7	\pm 12	\pm 7	\pm 5	\pm 6	\pm 7	\pm 9
			Change											
Timolol Maleate (Timoptic) 59 μ l of 0.5% Solution	4	106 \pm 6	-2	-4	-3	-5	-3	-8	-5	-3	-1	-6	-2	-5
			\pm 6	\pm 6	\pm 7	\pm 7	\pm 7	\pm 26	\pm 16	\pm 9	\pm 8	\pm 5	\pm 9	\pm 6
			Change											
Timolol Maleate (Timoptic) 59 μ l of 0.5% Solution	4	106 \pm 6	-2	-4	-3	-5	-3	-8	-5	-3	-1	-6	-2	-5
			\pm 6	\pm 6	\pm 7	\pm 7	\pm 7	\pm 26	\pm 16	\pm 9	\pm 8	\pm 5	\pm 9	\pm 6
			Change											

^aAll values are expressed as the mean \pm 1 S.D.

^bp < 0.05

TABLE VI
THE EFFECT OF TOPICALLY ADMINISTERED CELIPROLOL AND TIMOLOL ON HEART RATE (HRP) IN ANESTHETIZED DOGS

Treatment	N	Baseline ^a Pre-treatment (mm Hg)	Time (Minutes Posttreatment)											
			15	30	45	60	90	120	150	180	210	240	270	300
Saline 50 μ l 0.9% NaCl Solution	4	122 +24	Change (bpm)	6 +11	5 +20	3 +15	0 +20	0 +14	8 +10	-9 +26	-3 +20	-5 +25	-13 +31	-12 +18
		\bar{x}	Change	6 +11	3 +15	3 +12	1 +16	0 +11	7 +9	-7 +18	-1 +15	-3 +19	-9 +23	-10 +14
Celiprolol HCl (RUC 5320-A) 50 μ l 0.5% Solution	4	137 +21	Change (bpm)	-5 +6	-10 +10	-8 +18	-5 +15	-8 +11	-5 +18	-21 +20	-11 +16	-11 +9	-18 +23	-28 +10
		\bar{x}	Change	-4 +4	-8 +9	-6 +15	-4 +11	-5 +9	-5 +14	-17 +15	-9 +12	-9 +8	-14 +17	-22 +10
Timolol Maleate (Timoptic) 50 μ l of 0.5% Solution	4	119 +22	Change (bpm)	-12 +11	-19 +11	-25 ^b +17	-28 +18	-25 +25	-17 +11	-19 +38	-28 +33	-23 +37	-26 +32	-29 +26
		\bar{x}	Change	-9 +9	-15 +8	-19 ^b +12	-22 +13	-18 +21	-10 +39	-11 +35	-19 +29	-15 +33	-18 +29	-22 +22

^aAll values are expressed as the mean \pm 1 S.D.
^bp < 0.05

1 In asthmatic patients airway constriction has been
reported as the result of the use of certain β -blockers. The
present invention, utilizing celiprolol, provides for the
treatment of glaucoma without having the deleterious side
5 effect of airway constriction associated with certain
 β -blockers, such as atenolol, metoprolol, propanolol and
timolol. Example III describes the comparative study
conducted on celiprolol with three other beta blocking agents
for effects on bronchomotor tone in mechanically ventilated
10 cats infused with serotonin.

15

20

25

30

35

EXAMPLE IIIMETHOD

Male cats (3-5 kg) were anesthetized with pentobarbital sodium, 35 mg/kg i.p. Maintenance anesthetic was administered intravenously as needed. The trachea of each cat was cannulated and pneumotachograph placed on-line for monitoring air flow. A differential pressure transducer placed between the tracheal cannula and a cannula in the pleural cavity was used to monitor transpulmonary pressure. Electronic signals proportional to air flow and transpulmonary pressure were converted by an on-line analog computer to values of pulmonary (airway) resistance, R_{AW} , for each breath. Cannulae were inserted into the right femoral artery and both femoral veins. The arterial cannula was used for monitoring blood pressure and heart rate. In addition, it allowed for withdrawal of arterial blood samples for assessment of PO_2 , PCO_2 and pH, as a means of confirming the adequacy of ventilation. The animal was paralyzed with gallamine triethiodide (Flaxedil, 20 mg i.v.) and mechanically ventilated.

In order to test for bronchodilator activity, it was necessary to increase the normally low bronchomotor tone of the cat. This increase was induced by a constant i.v. infusion of serotonin (5-HT), approximately 20 μ g/kg/min. During this steady state bronchoconstriction, each animal received three or four increasing bolus doses of a single beta blocker, either atenolol (1-10 mg/kg i.v.), celiprolol hydrochloride (1-10 mg/kg i.v. or 1-100 mg/kg i.d.), metoprolol tartrate (1-10 mg/kg i.v.) or timolol maleate (0.03-3 mg/kg i.v.). Doses are expressed as the free bases. All drugs were dissolved in 0.9% saline, except timolol, which was used in the form of Timoptic [®] ophthalmic

1 solution suitably diluted with saline. In experiments in which celiprolol was given intraduodenally, needle-tipped cannulae were surgically inserted into the duodena of cats fasted for 16 hours.

5 Drug-induced changes in bronchomotor tone, averaged over 6-second intervals, were calculated as percent changes in R_{AW} from the steady state values established by serotonin infusion. Mean \pm standard deviation were calculated for experiments in which $n = 3$. Data for one celiprolol dose, 10 mg/kg i.d., were calculated using $n = 2$.
10 Statistical analysis was carried out with the t-test for paired data.

RESULTS

The results are shown in Table VII wherein R_{AW} =
15 % change in airway resistance; HR = heart rate - breaths/minute; i.v. = intravenous; i.d. = intraduodenal.

TABLE VII

20	<u>COMPOUND</u>	<u>mg/kg i.v.</u>	<u>HR</u>	<u>RAW</u>
	ATENOLOL	1	-8+5	23+12
	"	3	-6+3	14+5
	"	10	-6+2	48+34
25	METOPROLOL	1	-17+11	50+23
	"	3	-25+19	43+21
	"	10	-94+86	105+37
	TIMOLOL	0.3	-8+8	11+9
	"	1	-8+1	18+51
30	CELIPROLOL	1	-1+10	-62+10
	"	3	-9+4	-65+10
	"	10	-27+7	-70+14

1 As shown in the table, intravenous administration
of atenolol, metoprolol and timolol caused dose-dependent
bronchoconstriction, i.e., increased R_{AW} . Bronchoconstriction
has increased to 48% with atenolol, to 105% with
5 metoprolol and to 49% with timolol. In contrast, celiprolol
produced bronchodilation, i.e., R_{AW} decreased to -70%.

The duration of the broncodilator effect of
celiprolol was about 14 minutes at 1 and 3 mg/kg. The effect
of 10 mg/kg lasted for more than 40 minutes.

10

15

20

25

30

35

EXAMPLE IV

1 The procedure in this example generally follows the
procedure used in Example III, except that the test compounds
of celiprolol and timolol were administered topically as
5 follows. 50 μ l of test solution was instilled into the eyes
of anesthetized cats whose airway resistance, R_{AW} , had been
increased by infusion of serotonin. Three doses of timolol
(5 mg/ml) further increased R_{AW} by 20 to 35%. Similar
doses of celiprolol decreased R_{AW} by 21 to 23%.

10

15

20

25

30

35

EXAMPLE V

1 This procedure in this sample generally follows the
procedure used in Example III, except that celiprolol was
administered to cats intraduodenally. Tested in the range of
5 10-100 mg/kg, celiprolol; reduced R_{AW} by 29 to 41%.

 Having described the invention, those skilled in
the art will know modifications within the spirit thereof,
and the invention is to be limited only within the scope of
the appended claims.

10

15

20

25

30

35

1 WHAT IS CLAIMED IS:

1. An ophthalmic preparation for the treatment of glaucoma comprising an effective amount of a celiprolol salt in a pharmaceutically acceptable ophthalmic carrier for lowering intraocular pressure.
- 5 2. The ophthalmic preparation of Claim 1 wherein the celiprolol salt is present from about 0.1 to about 5.0% w/v concentrations in said pharmaceutically acceptable carrier.
- 10 3. The ophthalmic preparation of Claim 1 or 2 wherein from about 0.03 to about 0.5% w/v of the celiprolol salt is present in a pharmaceutically acceptable carrier.
4. The ophthalmic preparation of Claim 2 or 3 wherein said pharmaceutically acceptable carrier comprises in
- 15 % w/v:
- | | |
|---|------|
| boric acid-sodium borate | 0.03 |
| benzalconium chloride | 0.01 |
| disodium ethylenediamine | |
| tetraacetate | 0.1 |
| sodium thiosulfite | 0.3 |
| 20 polyvinylpyrrolidone | 1.5 |
| hydroxyethyl cellulose | 0.1 |
| polysorbate 80 | 0.01 |
| water QS to 100, and having an isotonicity range of | |
| 25 270 to 330 milliosmoles. | |
| 5. The ophthalmic preparation of any of Claims 1 to 4 wherein said pharmaceutically acceptable carrier is a water base or an ointment base. | |
| 6. The ophthalmic preparation of any of Claims 1 to 5 wherein the isotonicity of said preparation is within | |
| 30 the range of 270 to 330 milliosmoles. | |

1 7. The ophthalmic preparation of any of Claims 1
to 6 wherein the pH of said preparation is in the range of 6
to 9.0.

5 8. The ophthalmic preparation of any of Claims 1
to 7 wherein said celiprolol salt is celiprolol
hydrochloride.

10 9. The ophthalmic preparation of any of Claims 1
to 8 further comprising a buffer, a tonicity agent, a
preservative, a stabilizer, a co-solvent and a viscosity
agent.

10 10. An ophthalmic preparation for the treatment of
glaucoma according to Claim 1 comprising:

a. celiprolol hydrochloride 0.03-2.0 w/v;

15 b. a co-solvent selected from the group consisting
of glycerin, propylene glycol, polyethylene glycol,
polysorbates 20, 60 and 80 or Pluronic F-68;

20 c. a viscosity agent selected from the group
consisting of polyvinylpyrrolidone, polyvinyl alcohol, methyl
cellulose, hydroxypropyl methylcellulose, hydroxyethyl
cellulose and carboxymethylcellulose; sodium carboxymethyl-
cellulose or hydroxypropylcellulose;

25 d. a stabilizer selected from the group consisting
of sodium bisulfite, sodium thiosulfite, cysteine, acetyl
cysteine, β -cyclodextrin, dextran and thiourea,
thio-sorbitol, moroethioglycerol, sodium EDTA, or sodium
sulfosuccinate;

30 e. a 0.01 to 0.20 molar buffer selected from the
group consisting of boric acid-sodium borate, phosphate
buffer, boric acid-sodium bicarbonate, boric acid-sodium
citrate, citric acid-sodium phosphate, tri(hydroxymethyl)
amino methane-maleic acid and tris(hydroxymethyl) amino
methane-HCl;

1 f. a preservative selected from the group
consisting of benzalkonium chloride, disodium ethylene-
diamine tetracetate, thimerosal, chlorobutanol,
phenylmercuric nitrate, phenylmercuric acetate, methyl
5 paraben, propyl paraben, phenyl mercuric borate or
phenylethyl alcohol; and

g. water Q.S. to 100%.

11. A process of preparing an ophthalmic
preparation for lowering intraocular pressure comprising:

10 a. dissolving or dispersing an effective amount of
celiprolol salt in a pharmaceutically acceptable carrier;

b. adding buffer salts, stabilizers or viscosity
agents;

15 c. further optionally adding sodium chloride to
adjust to the required isotonicity;

d. adjusting to the final volume with purified
water.

12. The process according to Claim 1 wherein the
preparation is further sterilized by filtration or
20 autoclaved, or both.



European Patent
Office

EUROPEAN SEARCH REPORT

0109561

Application number

EP 83 11 0477

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
D, X	DE-A-2 458 624 (LENTIA) * Claims; example 1 * -----	1-12	A 61 K 31/17
			TECHNICAL FIELDS SEARCHED (Int. Cl. 7)
			A 61 K 31/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 02-02-1984	Examiner GAUTIER R.H.A.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	